

¹⁴ The author is indebted for technical assistance in the preparation of these matings to W. Gencarella.

¹⁵ No definite case of a double crossover in the $y - spl$ region was observed.

¹⁶ Two verified crossovers between y and sc were detected as $y^2 sc w spl$ phenotypes.

¹⁷ Morgan, T. H., Bridges, C. B., and Schultz, J., *Yearbook Carnegie Inst.*, **30**, 408-415 (1931).

¹⁸ Panshin, I. B., *Compt. rend. acad. sci. U. R. S. S.*, **30**, 57-60 (1941).

¹⁹ Bridges, C. B., *J. Heredity*, **29**, 11-13 (1938).

²⁰ Other studies cited by Bridges and Brehme,⁸ indicating that this break lies between 3C2 and 3C3 would not alter the present argument.

²¹ Horowitz, N. H., these PROCEEDINGS, **31**, 153-157 (1945).

ON THE NUMBER OF BOUND STATES IN A CENTRAL FIELD OF FORCE

BY V. BARGMANN

PRINCETON UNIVERSITY

Communicated by E. P. Wigner, September 18, 1952

1. The present note contains some fairly elementary remarks concerning the number of bound state solutions of the Schrödinger equation

$$\nabla^2\psi + E\psi = V(r)\psi$$

for a central field of force, more specifically, the number n_l of bound state solutions of the radial wave equation

$$\phi'' - l(l+1)r^{-2}\phi + E\phi = V(r)\phi \quad (1)$$

for angular momentum l . We assume the integral

$$I = \int_0^\infty r |V(r)| dr \quad (2)$$

to be *finite*, and we wish to estimate n_l in terms of I . (In the units chosen V has the dimension (length)⁻², so that I is dimensionless.) R. Jost and A. Pais (ref. 1, p. 844) have shown that no bound states occur if $I < 1$. Our aim is to derive the more general inequality

$$(2l+1)n_l < I \quad (3)$$

(equality excluded). The number n_l counts the distinct stationary energy values corresponding to equation (1). If the $(2l+1)$ -fold degeneracy of each of them is taken into account it is seen that for a given angular momentum l there are less than I bound states, and no bound states occur if $l \geq \frac{1}{2}(I-1)$. The estimate (3) is best possible in the sense that for a given l potentials may be constructed which have a prescribed number n_l of bound states for that angular momentum and for which I approaches

$(2l + 1)n_i$ arbitrarily closely (see section 5 below). The whole question is taken up because the finiteness of I plays a significant role in several recent investigations on scattering theory.¹⁻⁴ (V may have any singularities consistent with a finite value of I .)

2. As is well known, n_i is the number of zeros (not counting $r = 0$) of that solution $\phi(r)$ of the equation

$$\phi'' - l(l + 1)r^{-2}\phi = V(r)\phi \quad (4)$$

($E = 0$) which vanishes at the origin. Special care must be taken with a possible bound state $E = 0$. Since I is finite any solution of (4) has the following asymptotic behavior at infinity. The expression $r^{-(l+1)}\phi(r)$ always approaches a finite limit, say λ , as $r \rightarrow \infty$. If $\lambda \neq 0$, $\phi(r)$ increases indefinitely. If $\lambda = 0$ the expression $r^l\phi(r)$ approaches a finite limit μ , and $\mu \neq 0$. In the latter case $\phi(r)$ is square integrable if $l > 0$, and accordingly $E = 0$ is a bound state.⁵ For $l = 0$, $E = 0$ is never a bound state if I is finite. For the purpose of our discussion, however, we shall count $r = \infty$ as a zero of $\phi(r)$ —even if $l = 0$ —whenever $\lim_{r \rightarrow \infty} r^{-(l+1)}\phi(r) = 0$, and interpret the inequality (3) accordingly.⁶

Replace in equation (1) $V(r)$ by a potential $V_1(r)$ such that $V_1(r) \leq V(r)$ for all r , and denote by n_i' the number of bound states for the new potential. Then $n_i' \geq n_i$. We shall choose $V_1(r) = -W(r)$, where $W(r) = |V(r)|$, and study the equation

$$\phi'' - l(l + 1)r^{-2}\phi = -W(r)\phi \quad W(r) = |V(r)|. \quad (5)$$

Denote by $\nu_1, \nu_2, \dots, \nu_n$ ($n = n_i'$) the successive zeros of $\phi(r)$ ($0 < \nu_1 < \nu_2 < \dots < \nu_n$), and set $\nu_0 = 0$. We shall prove

$$\int_{\alpha}^{\beta} r W(r) dr > 2l + 1; \quad \alpha = \nu_{k-1}, \beta = \nu_k, k \geq 1. \quad (6)$$

The inequality (3) is obtained by adding the n inequalities (6), for we find then

$$I = \int_0^{\infty} r W(r) dr \geq \int_0^{\nu_n} r W(r) dr > n_i'(2l + 1) \geq n_i(2l + 1).$$

3. *Preliminary Remarks on $\phi(r)$.*—The solution of equation (5) which vanishes at the origin is uniquely determined up to a constant factor. As $r \rightarrow 0$, $r^{-(l+1)}\phi(r)$ approaches a finite non-vanishing limit κ . Choosing $\kappa = 1$, we find from (5)

$$\phi(r) = r^{l+1} - \int_0^r G(r, \rho)\phi(\rho)W(\rho) d\rho. \quad (7)$$

$\phi(r)$ is then real. Here $G(r, \rho)$ is the fundamental solution of the equation $f'' - l(l + 1)r^{-2}f = 0$, i.e., $\partial^2 G(r, \rho)/\partial r^2 - l(l + 1)r^{-2}G(r, \rho) = 0$, $G(r, r) = 0$, and $\partial G(r, \rho)/\partial r = 1$ for $r = \rho$. We have

$$G(r, \rho) = (2l+1)^{-1}H(r, \rho); \quad H(r, \rho) = r(r/\rho)^l - \rho(\rho/r)^l \quad (8)$$

if $r > 0, \rho > 0$. Clearly $H(r, \rho) > 0$ if $r > \rho$. Since, for $r \rightarrow 0, r^{-(l+1)}\phi(r) \rightarrow 1$, the integral in (7) is absolutely convergent.

In the sequel we shall need the inequality

$$H(\beta, \rho)H(\rho, \alpha) \leq \rho(H(\beta, \alpha) - Y(\beta, \alpha))$$

$$Y(\beta, \alpha) = 2(\alpha\beta)^{1/2}[1 - (\alpha/\beta)^{l+1/2}] > 0 \quad (\beta > \alpha). \quad (9)$$

To derive it consider $Z(\rho, \beta, \alpha) = \rho H(\beta, \alpha) - H(\beta, \rho)H(\rho, \alpha)$. By straight forward computation

$$\begin{aligned} Z(\rho, \beta, \alpha) &= \rho(\alpha\beta)^{1/2}\{(\rho^2/\alpha\beta)^{l+1/2} + (\alpha\beta/\rho^2)^{l+1/2} - 2(\alpha/\beta)^{l+1/2}\} \\ &= \rho(\alpha\beta)^{1/2}\{[(\rho/\sqrt{\alpha\beta})^{l+1/2} - (\sqrt{\alpha\beta}/\rho)^{l+1/2}]^2 + 2[1 - (\alpha/\beta)^{l+1/2}]\} \\ &\geq \rho Y(\beta, \alpha) \end{aligned}$$

which establishes (9).

4. *Proof of (6).*—We distinguish four cases' according as $\alpha = 0, \alpha > 0; \beta < \infty, \beta = \infty$.

(a) $\alpha = 0, \beta = \nu_1 < \infty$. On the open interval $(0, \beta)$ ϕ is positive, and hence, by (7), $\phi(r) \leq r^{l+1}$. Since $\phi(\beta) = 0$ we have from (7) and (8)

$$\begin{aligned} (2l+1)\beta^{l+1} &= \int_0^\beta H(\beta, \rho)\phi(\rho)W(\rho)d\rho \leq \int_0^\beta H(\beta, \rho)\rho^{l+1}W(\rho)d\rho \\ (2l+1)\beta^{l+1} &\leq \beta^{l+1}[\int_0^\beta \rho W(\rho)d\rho - \int_0^\beta (\rho/\beta)^{2l+1}\rho W(\rho)d\rho]. \end{aligned}$$

On dividing by β^{l+1} we find (6) because the last integral is positive.

(b) $\alpha > 0, \beta < \infty$. Since $\phi(\alpha) = 0$, the derivative $\phi'(\alpha)$ does not vanish. If we replace $\phi(r)$ by $\chi(r) = \phi(r)/\phi'(\alpha)$, then $\chi(\alpha) = 0, \chi'(\alpha) = 1$, and hence

$$\chi(r) = G(r, \alpha) - \int_\alpha^r G(r, \rho)\chi(\rho)W(\rho)d\rho \quad (10)$$

On the interval $\alpha < r < \beta$, therefore, $0 < \chi(r) \leq G(r, \alpha)$. Thus, for $r = \beta$,

$$G(\beta, \alpha) = \int_\alpha^\beta G(\beta, \rho)\chi(\rho)W(\rho)d\rho \leq \int_\alpha^\beta G(\beta, \rho)G(\rho, \alpha)W(\rho)d\rho,$$

or

$$\begin{aligned} (2l+1)H(\beta, \alpha) &\leq \int_\alpha^\beta H(\beta, \rho)H(\rho, \alpha)W(\rho)d\rho \\ &\leq [H(\beta, \alpha) - Y(\beta, \alpha)] \int_\alpha^\beta \rho W(\rho)d\rho \end{aligned}$$

[see (9)]. Division by $H(\beta, \alpha)$ establishes (6).

(c) $\alpha = 0, \beta = \nu_1 = \infty$. Here we use (7), and, as in case (a), $r^{l+1} \geq \phi(r) > 0$ for all positive r . By assumption, $\tau(r) = (2l+1)r^{-(l+1)}\phi(r)$ approaches 0 as $r \rightarrow \infty$. By (7),

$$2l+1 - \tau(r) - \int^r K(r, \rho)\phi(\rho)W(\rho)d\rho = 0$$

where

$$K(r, \rho) = r^{-(l+1)} H(r, \rho) = \rho^{-l} (1 - (\rho/r)^{2l+1}) < \rho^{-l}$$

Hence

$$\int_0^r \rho W(\rho) d\rho \geq \int_0^r \rho^{l+1} K(r, \rho) W(\rho) d\rho \geq \frac{2l+1-\tau(r)}{2l+1-\tau(r) + \int_0^r K(r, \rho) [\rho^{l+1} - \phi(\rho)] W(\rho) d\rho} \quad (11)$$

Since the integral in (11) is non-negative and $\tau(r) \rightarrow 0$, we find at once that $\int_0^\infty \rho W(\rho) d\rho \geq 2l+1$. To exclude equality we observe that there must exist two adjacent intervals $[\xi, \eta]$ and $[\eta, \zeta]$ ($\xi < \eta < \zeta < \infty$) such that $\int_\xi^\eta \rho W(\rho) d\rho$ and $\int_\eta^\zeta \rho W(\rho) d\rho$ both exceed $1/4$, say. If $r \geq \eta$, then, by (7), $r^{l+1} - \phi(r) \geq \int_\xi^\eta G(r, \rho) \phi(\rho) W(\rho) d\rho$ and throughout the interval $[\eta, \zeta]$ $(\rho^{l+1} - \phi(\rho)) \rho^{-(l+1)} \geq c$, where c is some positive constant. For $r > \zeta$ we find from (11)

$$\int_0^r \rho W(\rho) d\rho \geq 2l+1 - \tau(r) + c \int_\eta^\zeta \rho^l K(r, \rho) \rho W(\rho) d\rho$$

and in the limit $r \rightarrow \infty$

$$\int_0^\infty \rho W(\rho) d\rho \geq 2l+1 + c \int_\eta^\zeta \rho W(\rho) d\rho > 2l+1, \quad \text{q. e. d.}$$

(d) $\alpha > 0$, $\beta = \nu_n = \infty$. We start, as in (b), from equation 10, so that $G(r, \alpha) \geq \chi(r) > 0$ for $r > \alpha$. By assumption, $(2l+1)^{-1} \theta(r) = \chi(r)/G(r, \alpha)$ approaches 0 as $r \rightarrow \infty$. From (10) we find

$$2l+1 - \theta(r) - \int_\alpha^r \frac{H(r, \rho)}{G(r, \alpha)} \chi(\rho) W(\rho) d\rho = 0.$$

Hence, by (9),

$$\int_\alpha^r \rho W(\rho) d\rho \geq \int_\alpha^r \frac{H(r, \rho)}{G(r, \alpha)} G(\rho, \alpha) W(\rho) d\rho \geq 2l+1 - \theta(r) + \int_\alpha^r \frac{H(r, \rho)}{G(r, \alpha)} [G(\rho, \alpha) - \chi(\rho)] W(\rho) d\rho \quad (12)$$

We proceed as in case (c) above. The inequality $\int_\alpha^\infty \rho W(\rho) d\rho \geq 2l+1$ is an immediate consequence of (12). To exclude equality the intervals $[\xi, \eta]$ and $[\eta, \zeta]$ are chosen as before ($\xi \geq \alpha$), so that $(2l+1)(G(\rho, \alpha) - \chi(\rho)) \rho^{-(l+1)} \geq c' > 0$ if $\eta \leq \rho \leq \zeta$. For $r > \zeta$ (12) implies

$$\int_\alpha^r \rho W(\rho) d\rho \geq 2l+1 - \theta(r) + c' \int_\eta^\zeta \rho^l \frac{H(r, \rho)}{H(r, \alpha)} \rho W(\rho) d\rho$$

and since $\lim_{r \rightarrow \infty} (H(r, \rho)/H(r, \alpha)) = (\alpha/\rho)^l$, we find for $r \rightarrow \infty$ $\int_\alpha^\infty \rho W(\rho) d\rho \geq 2l+1 + c' \alpha^l \int_\eta^\zeta \rho W(\rho) d\rho > 2l+1$. This concludes the proof of (6).

5. *Examples.*—The proofs in the preceding section suggest the construction of potentials for which the inequalities (3) or (6) may be approxi-

mately replaced by the corresponding equalities. In 4(a), for example, the first inequality will nearly reduce to an equality if at those r where $W(r)$ is appreciable $\phi(r)$ nearly equals r^{l+1} , i.e., the field free solution. This leads (for $n_l = 1$) to the choice of a potential $V(r) = -W(r)$ ($W \geq 0$) which vanishes everywhere with the exception of a small interval $a < r < a + \delta = b$. Outside $[a, b]$ we obtain for a suitably normalized solution of equation (5)

$$\left. \begin{aligned} \phi(r) &= (r/a)^{l+1} & r \leq a \\ \phi(r) &= c_1(b/r)^l - c_2(r/b)^{l+1} & r \geq b \end{aligned} \right\} \quad (13)$$

$$\left. \begin{aligned} (2l+1)c_1 &= (l+1)\phi(b) - b\phi'(b); \\ (2l+1)c_2 &= -l\phi(b) - b\phi'(b) \end{aligned} \right\} \quad (14)$$

If $c_2 > 0$, $\phi(r) \rightarrow -\infty$ as $r \rightarrow \infty$, so that $\phi(r)$ vanishes at a point β given by $(\beta/b)^{2l+1} = c_1/c_2$, and if $c_2 = 0$, then $\beta = \infty$. Owing to the smallness of δ the relative change of $\phi(r)$ across the interval $[a, b]$ is negligible compared to the relative change of $\phi'(r)$, so that $\phi(b) \sim \phi(a)$. From the condition $c_2 \geq 0$ we obtain $-\phi'(b)/\phi(b) > l/b$, and since $\phi'(a)/\phi(a) = (l+1)/a$, this amounts to

$$\phi'(a)/\phi(a) - \phi'(b)/\phi(b) \sim (\phi'(a) - \phi'(b))/\phi(a) \gtrsim (2l+1)/a \quad (15)$$

Thus the required increment of the logarithmic derivative is the smaller the larger a is chosen—or a potential of given strength is the more effective in producing bound states the farther it is removed from the origin (which is the reason for the weight factor r in the integral I). Without yet specifying $W(r)$, we see from (5) that $\phi'(b) - \phi'(a)$ approximately equals $-\bar{W}\delta\phi(a)$ where \bar{W} is a suitable average of W , provided the centrifugal term $l(l+1)r^{-2}$ is negligible compared to \bar{W} . If the increment is as small as possible we find from (15) that $\bar{W}\delta a \sim (2l+1)$ which is equivalent to $\int_a^b rW(r) dr = \int_0^\infty rW(r) dr \sim 2l+1$.

To have a definite example consider $W(r) = 1 + l(l+1)r^{-2}$, and $a = (2l+1)(1+\delta)/\delta$. Then, in $[a, b]$, $\phi(r) = \cos(r-a) + ((l+1)/a)\sin(r-a)$ so that $\phi(b) = \cos\delta + ((l+1)/a)\sin\delta$, $\phi'(b) = -\sin\delta + ((l+1)/a)\cos\delta$, and one verifies easily that $c_2 > 0$ for small δ (e.g., $\delta < 1/4$). The zero, β , is determined by $(\beta/b)^{2l+1} = c_1/c_2$, and approximately $c_1/c_2 \sim \delta^{-1}$ so that $\beta \sim a \cdot \delta^{-1/(2l+1)}$. Finally, $I = \int_a^b rW(r) dr = \delta(a + 1/2\delta) + l(l+1)\log(1+\delta/a)$. As $\delta \rightarrow 0$, $I \rightarrow 2l+1$. Alternately, instead of varying a and keeping the strength of W fixed, one might keep a fixed and vary the strength of the potential.

In a similar way one may construct potentials with two or more bound state solutions such that I is arbitrarily close to $(2l+1)n_l$. One simply has to add other troughs in suitably placed intervals $[a', a' + \delta']$, etc., in such a way, however, that two successive intervals are sufficiently far

from one another and from the zero of $\phi(r)$ between them. Note that these potentials are adjusted only to one fixed value of the angular momentum.

¹ Jost, R., and Pais, A., *Phys. Rev.*, **82**, 840 (1951).

² Levinson, N., *Kgl. Danske Videnskab. Selskab. Mat.-fys. Medd.*, **25**, No. 9 (1949).

³ Jost, R., and Kohn, W., *Phys. Rev.*, **87**, 977 (1952).

⁴ Bargmann, V., *Rev. Mod. Phys.*, **21**, 488 (1949).

⁵ For $l = 0$ these statements are proved in the appendix of ref. 4. For higher l a similar proof may be given.

⁶ Even for $l = 0$ this case is significant although no bound state is present. In particular in this case $|\sin \eta(0)| = 1$ where $\eta(k)$ is the scattering phase shift, and hence the cross section $\sin^2 \eta(k)/k^2$ becomes infinite as $k \rightarrow 0$. (See ref. 4, equation (1.9). In this case $f(0) = 0$.)

⁷ For $l = 0$ one need not distinguish $\alpha = 0$ and $\alpha > 0$, because $G(r, \alpha) = r - \alpha$ may be used in both cases.

ON THE INVARIANT THEORY OF THE CLASSICAL GROUPS

By F. D. MURNAGHAN

INSTITUTO TECNOLÓGICO DE AERONÁUTICA, SÃO JOSÉ DOS CAMPOS, BRASIL

Communicated August 18, 1952

It has been recognized for some time that the theory of invariants and covariants, with respect to a given group, rests on the analysis of $\Gamma \otimes \Gamma'$, where Γ and Γ' are irreducible representations of the group, into its irreducible components. Thus if we denote by $\{\lambda\}$ the irreducible representation of the n -dimensional linear group which is associated with the partition $(\lambda) = (\lambda_1, \dots, \lambda_n)$, $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n \geq 0$, of any non-negative integer m into not more than n parts the core of the theory of invariants and covariants, under linear transformations, is the analysis of $\{\lambda\} \otimes \{\mu\}$ where (λ) and (μ) are partitions of any two non-negative integers m and j , respectively. The cases where (λ) is either the 1-element partition (m) or the m -element partition (1^m) and (μ) is either the 1-element partition (j) or the j -element partition (1^j) are of particular importance and the problem of analyzing $\{\lambda\} \otimes \{\mu\}$, especially in these cases, has been much studied, following the initial impetus given by Littlewood,¹ during the past decade. However the methods used have been laborious when m and j are greater than 2; in these cases $\{\lambda\} \otimes \{\mu\}$ contains many components, each corresponding to a partition of mj , and each of these has had to be determined separately by a tedious calculation. We present in this note a method which yields, in the cases of particular importance referred to, the components of $\{\lambda\} \otimes \{\mu\}$ in platoons, rather than individually, each platoon consisting of those parentheses $\{\dots\}$ which contain the same number of non-zero parts.